

A supplement to my paper on real and complex operator ideals

Jörg Wenzel

February 8, 2008

Abstract

We show that the main problem left open in [2] can be solved using the Banach spaces Z_α recently constructed by Kalton [1]. This gives an example of a complex operator ideal that has no real analogue. It thus shows the richer structure of complex operator ideals compared with the real ones.

1 Introduction

For a Banach space X let $\overline{X}$ denote its complex conjugate, i.e. the Banach space X equipped with the scalar multiplication $\alpha \odot x := \overline{\alpha}x$.

For an operator $T : X \rightarrow Y$ let $\overline{T}$ denote its complex conjugate, i.e. the operator T acting between $\overline{X}$ and $\overline{Y}$. This is again a linear operator.

For a complex operator ideal $\mathfrak{C}$ the complex conjugate ideal $\overline{\mathfrak{C}}$ is defined as the ideal consisting of all T such that $\overline{T} \in \mathfrak{C}(\overline{X}, \overline{Y})$.

The ideal $\mathfrak{C}$ is called self conjugate if $\mathfrak{C} = \overline{\mathfrak{C}}$.

The problem of the existence of operator ideals that are not self conjugate was left open in [2]. However, it was reduced there to the problem of finding a complex Banach space that is not isomorphic to its complex conjugate, but that is isomorphic to its square.

In [1], Kalton gives examples of Banach spaces Z_α which are not isomorphic to their complex conjugates. On the other hand, the elementary nature of these spaces makes it possible to easily verify that they are isomorphic to their squares, thus yielding the required Banach spaces.

Research supported by German Academic Exchange Service (DAAD)
1991 *Mathematics Subject Classification.* 47D50, 46B20.

2 The spaces Z_α

In [1] Kalton constructed elementary examples of Banach spaces Z_α , not isomorphic to their complex conjugates. We need only one additional property of these spaces which is not explicitly mentioned in [1], namely that Z_α is Cartesian. This is however easy, as is shown below.

For the convenience of the reader, let us repeat here the definition of Z_α .

Let $f_\alpha(t) := t^{1+i\alpha}$ for $-\infty < \alpha < \infty$ and $0 \leq t < \infty$. Given a sequence $x = (\xi_k) \in l_2$ define the sequence $\Omega_\alpha(x)$ by

$$(\Omega_\alpha(x))_k := \begin{cases} \xi_k f_\alpha\left(\log \frac{\|x\|_2}{|\xi_k|}\right) & \text{if } \xi_k \neq 0 \\ 0 & \text{otherwise.} \end{cases}$$

Let Z_α be the space of all pairs of complex-valued sequences (x, y) such that

$$\|(x, y)\|_\alpha := \|x\|_2 + \|y - \Omega_\alpha(x)\|_2 < \infty.$$

It turns out that Z_α is a Banach space under a norm equivalent to the quasi norm $\|\cdot\|_\alpha$.

Moreover $\overline{Z_\alpha} = Z_{-\alpha}$ and Z_α is isomorphic to Z_β if and only if $\alpha = \beta$.

Definition. A Banach space X is called *Cartesian* if it is isomorphic to its Cartesian square $X \oplus X$.

Proposition. *The spaces Z_α are Cartesian.*

PROOF: For a complex-valued sequence $x = (\xi_k)$ define

$$U^{odd}x := (\xi_{2k-1}) \quad \text{and} \quad U^{even}x := (\xi_{2k}).$$

Then the map $U : Z_\alpha \rightarrow Z_\alpha \oplus Z_\alpha$ defined by

$$U(x, y) := (U^{odd}x, U^{odd}y) \oplus (U^{even}x, U^{even}y)$$

defines an isomorphism.

The bijectivity of U is trivial, so we only show its continuity.

It is mentioned in [1] that there is a constant C , such that for $s \in l_\infty$ we have the estimate

$$\|(sx, sy)\|_\alpha \leq C \cdot \|s\|_\infty \cdot \|(x, y)\|_\alpha.$$

Hence

$$\|(U^{odd}x, U^{odd}y)\|_\alpha \leq C \|(x, y)\|_\alpha \quad \text{and} \quad \|(U^{even}x, U^{even}y)\|_\alpha \leq C \|(x, y)\|_\alpha.$$

This proves the assertion. $\square$

3 Main theorem

Theorem. *For $\alpha \neq 0$ the operator ideal*

$$\mathfrak{C}_\alpha := \{T : T \text{ admits a factorization over the space } Z_\alpha\}$$

is not self conjugate.

PROOF: Since Z_α is Cartesian, $\mathfrak{C}_\alpha$ is indeed an operator ideal. Now $Id_{Z_\alpha} \in \mathfrak{C}_\alpha$. Assume that $\overline{Id_{Z_\alpha}} = Id_{\overline{Z_\alpha}} \in \mathfrak{C}_\alpha$. This implies that Z_α and $\overline{Z_\alpha}$ are complemented in each other. Since Z_α and $\overline{Z_\alpha}$ are Cartesian, we can apply Pełczyński's decomposition method to obtain that Z_α and $\overline{Z_\alpha}$ are actually isomorphic which contradicts the properties of these spaces. See [2] for details. $\square$

References

- [1] KALTON, N. J. An elementary example of a Banach space not isomorphic to its complex conjugate. *Canad. Math. Bull.* **38** no. 2 (1995), 218–222.
- [2] WENZEL, J. Real and complex operator ideals. *Quaestiones Mathematicae* **18** no. 1–3 (1995), 271–285.

MATHEMATISCHES INSTITUT, FSU JENA, 07740 JENA, GERMANY
E-mail: wenzel@minet.uni-jena.de